

# Perspectives

Recovery Strategies from the OR to Home

## In This Issue

**C**onsiderable advances have been made in the diagnosis and treatment of breast cancer, however, this challenging disease is still the most common form of cancer in women. In 2001, nearly 200,000 women were diagnosed with breast cancer and over 40,000 succumbed to the disease. Ms Fortenbaugh discusses the methods of evaluations of palpable and non-palpable lesions, and, with a confirmed diagnosis, the treatment strategies. While the treatment is often multimodal including surgery, radiation, and chemotherapy, Ms. Fortenbaugh's article focuses on surgical intervention, postoperative care, and breast reconstruction.

Inflammatory bowel disease (Crohn's Disease and Ulcerative Colitis) are both difficult disorders for patients and clinicians. Crohn's Disease (CD) and Ulcerative Colitis (UC) are chronic, medically incurable conditions. Although genetic, biochemical, and immunologic factors are recognized as features of both disorders, definitive causes remain elusive. UC and CD occur among all age groups but have a peak incidence in the second and third decades. This article on ulcerative colitis is first in a series that will cover the medical and surgical management of inflammatory bowel disease. The second article will focus on the medical and surgical management of Crohn's disease.

## Advisory Board

Cheryl Bressler, MSN, RN, CORLN  
Oncology Nurse Specialist, Oncology Memorial Hospital, Houston, TX,  
National Secretary SOHN

Lois Dixon, MSN, RN  
Adjunct Faculty, Trinity College of Nursing, Moline, IL  
Pulmonary Staff Nurse, Genesis Medical Center, Davenport, IA

Jan Foster RN, PhD(c), MSN, CCRN  
Asst. Professor for Adult Acute and Critical Care Nursing  
Houston Baptist University, TX  
Secretary/Treasurer, AACN Certification Corp.

Mikel Gray, PhD, CUNP, CCCN, FAAN  
Nurse Practitioner/Specialist, Associate Professor of Nursing,  
Clinical Assistant Professor of Urology, University of Virginia,  
Department of Urology, Charlottesville, VA, Past-president SUNA

Victoria-Base Smith, PhD, MSN, CRNA, CCRN  
Clinical Assistant Professor, Nurse Anesthesia,  
University of Cincinnati, OH

Mary Sieggreen, MSN, RN, CS, NP  
Nurse Practitioner, Vascular Surgery, Harper Hospital, Detroit, MI

Franklin A. Shaffer, EdD, DSc, RN  
Vice-president, Education and Professional Development,  
Executive Director, Cross Country University

## The Perioperative Management of Breast Cancer

By Cathy Fortenbaugh MSN, AOCN, CSN, BSC, C

**B**reast cancer is the most common type of cancer in American women.<sup>1</sup> In fact, a woman is diagnosed with breast cancer every three minutes.<sup>1</sup> It touches the lives of all of us in some way. Last year, 193,700 women were diagnosed with breast cancer and 40,600 died of their disease.<sup>2</sup> Better diagnostic techniques, such as digital mammography, will help more breast cancers to be detected early, when they are most treatable.

Surgery is the first part of a multimodal approach for treating breast cancer that includes radiation and chemotherapy. Managing breast cancer is a team effort. Nurses are a crucial part of this team. A multidisciplinary approach is optimal. Ideally, a woman will consult with various members of the team in the same visit. The team members consist of surgical oncologists, plastic surgeons, radiologists, pathologists, radiation oncologists, medical oncologists, nurses, social workers, pastoral care, and nutritionists.

### Detection and diagnosis

Many breast abnormalities are found by women themselves or their healthcare providers. Mammography detects abnormalities that are too small to be palpated. After discovery, the approach to evaluation depends on whether an abnormality is palpable or nonpalpable. Breast cancer cannot be diagnosed without a tissue sample that has been evaluated by a pathologist.

### Palpable lesions

If the abnormality is palpable, there are

several ways to obtain a tissue sample:

- fine-needle aspiration
- core-needle biopsies
- incisional or excisional biopsies

A fine-needle aspiration can be done in the surgeon's office. The surgeon inserts a needle with a syringe into the mass and aspirates some cells. The cells are then placed on a slide, fixative is applied, and the slide is sent to a cytology lab. This approach can identify breast cancer cells but cannot distinguish between invasive and noninvasive breast cancer. If the specimen is positive for breast cancer, further evaluation of the tumor needs to be done.

A core-needle biopsy is a procedure in which the surgeon uses a larger needle to obtain a core of tissue. The area is numbed prior to biopsy with a local anesthetic. There may not be enough tissue to evaluate hormone receptor status. A core-needle biopsy is most commonly done when the breast mass is large and treatment options are limited.

Incisional biopsy is done on a large breast mass where the entire area of abnormal tissue cannot be removed. The surgeon makes an incision and removes enough tissue to obtain an accurate diagnosis.

Excisional biopsy, also known as lumpectomy, is recommended in most cases, especially for women with early-stage breast cancer. The purpose of lumpectomy is to re-



Continued on page 4

# Management of Patients with Inflammatory Bowel Disease. *Part 1: Ulcerative colitis*

By Janice Colwell, RN, MS, CWOCN

**I**nflammatory bowel disease (IBD) refers to Crohn's disease (CD) and ulcerative colitis (UC). Both are disorders of unknown origin involving genetic and immunologic influences on the ability of the gastrointestinal (GI) tract to distinguish foreign objects from self and to down-regulate the mucosal immune response.<sup>1</sup> The causes(s) of UC and CD are unknown. The absence of appropriate animal models for chronic IBD has hampered progress in determining its pathogenesis.

CD and UC share many overlapping clinical and therapeutic features. Both are chronic, medically incurable conditions. Although genetic, biochemical, and immunologic factors are recognized as features of both disorders, definitive causes remain elusive. UC and CD occur among all age groups but have a peak incidence in the second and third decades.

CD and UC are more common in developed countries, such as Scandinavia, Europe, and North America, and uncommon in Asia, Africa, and South America. The incidence of UC has remained stable; however, the incidence of CD has increased steadily. IBD is generally more common in urban than rural areas. Men are slightly more likely to be affected by UC and women with CD.<sup>2</sup>

Overall, 15-20% of patients have a family history of IBD. The family tendency is primarily genetic rather than due to a shared environmental etiology.<sup>3</sup>

It is clear that IBD is not the result of a single gene defect. Various studies have led to a consensus hypothesis that, in genetically predisposed individuals, both environmental factors (infectious, luminal bacteria) and endogenous factors (barrier function, vascular supply) induce intestinal mucosal immune responses that are abnormal in either their activation or regulation.<sup>4</sup> The abnormal immune response allows persistent amplification of the tissue-damaging process, resulting in mucosal tissue damage, diarrhea, and other immune responses.

This article on ulcerative colitis is first in a series that will cover the medical and surgical management of inflammatory bowel disease. The second article will focus on the medical

and surgical management of Crohn's disease.

## Ulcerative colitis

UC is a mucosal disease that occurs in the colon. It begins in the rectal mucosa and extends to a variable proximal level. In one-quarter of patients with UC, disease is limited to the rectum. In another 25-50%, the rectum and sigmoid or descending colon are involved. About one-third of patients have inflammation that extends proximal to the splenic flexure or involves the entire colon. The appearance of colonic mucosa can vary, depending on the severity of disease. Mild inflammatory changes include fine granularity of the mucosa and pinpoint hemorrhage. Moderate changes include coarse granularity, spontaneous hemorrhage, and the exudation of pus. In severe UC, inflammation extends into the submucosa, causing the bowel wall to become tissue-paper thin with a significant risk of spontaneous perforation (toxic megacolon). Cancer of the colon is a long-term complication of UC. The cancer risk increases with the duration and extent of disease and appears to increase progressively after 8-10 years of disease. Patients with pancolitis are at greater risk than patients with left-sided disease. Surveillance is recommended every 2 years or 8-10 years after the onset of disease in patients with pancolitis.<sup>1,5</sup>

## Presentation

The symptoms of UC depend on the extent and severity of inflammation. Patients with disease limited to the rectum demonstrate bleeding, tenesmus, passage of mucopus, and variation in the number and forms of stools. Many patients with UC proctitis have constipation. The greater the extent of bowel involved, the more likely the patient is to suffer from diarrhea. As the severity of the disease progresses, the patient suffers from systemic symptoms, such as low-grade fever, malaise, night sweats, and arthralgias.<sup>1,6</sup> Extraintestinal problems may include cutaneous disorders (pyoderma gangrenosum), ocular disturbances (uveitis), and liver problems (sclerosing cholangitis).

## Diagnosis

The diagnostic work-up of patients suspected of having ulcerative colitis will include a colonic endoscopy. Continuous mucosal changes with a distinct upper boundary to adjacent, normal-looking mucosa are typical signs of UC. Air-contrast barium studies of the colon reveal diffuse, contiguous granularity, superficial ulceration, and absent haustrations in active UC. Endoscopic and radiologic studies in addition to a biopsy confirm the diagnosis. Serologic measurement may be useful in differentiating UC from CD; perinuclear antineutrophil antibodies (pANCA) are present in most patients with UC but in a minority of patients with CD.<sup>7</sup>

## Medical treatment

The treatment of UC depends on the mucosal extent and severity of disease. The mucosal extent is important, because topical rectal therapies can be used to reach the disease. The severity of UC is relevant to estimate the intensity of anti-inflammatory therapy and risk of complications.<sup>8</sup> The goal of therapy has been to improve signs and symptoms of the disease.

*Aminosalicilyc acid derivatives of sulfasalazine* are the primary therapies for mild or moderate UC. Sulfasalazine was developed to combine an antibiotic (sulfapyridine) with a salicylate (5-aminosalicylic acid, mesalamine) for delivery into colonic tissue.<sup>8</sup> Sulfasalazine is effective in a dose-dependent manner for treating and maintaining the remission of UC.<sup>1</sup> Because 5-aminosalicylic acid (5-ASA) is the major source of therapeutic benefits and sulfapyridine is responsible for most side effects, sulfa-free aminosalicilylates have been developed recently. They include sulfasalazine (Azulfidine), olsalazine (Dipentum), Asacol (delayed release), and Pentasa (sustained release). Mesalamine suppository and enema formulations are available for rectal administration.<sup>9</sup>

Sulfasalazine side effects include nausea, headache, and myalgias. Sulfasalazine competitively inhibits the intestinal absorption of folate and can produce a folate deficiency. One milligram of folate daily is recommended for patients on long-term therapy.

*Corticosteroids* remain the mainstay of acute therapy for moderate, severe, and fulminant UC.<sup>10</sup> Corticosteroids are indicated to induce remission in moderate to severe UC at doses of 40-60 mg daily.<sup>11</sup> Hydrocortisone enemas are useful for treating distal colonic symptoms. Corticosteroid therapy is limited by its adverse effects: cushingoid features, acne, excess facial hair, and high blood pressure, and long-term effects of bone disease, cataract formation, and steroid-induced diabetes.

*Immunomodulators*, such as azathioprine (AZA) and 6-mercaptopurine (6-MP), have

been used in the long-term treatment of UC for over 30 years. This class of medications acts as an immunosuppressant and interferes with DNA and RNA synthesis.<sup>12</sup> These drugs enable the reduction or elimination of corticosteroids; however, they take from 3-6 months to reach maximal therapeutic doses. Intravenous cyclosporine has benefited the short-term management of severe, refractory UC in 50-80% of patients.<sup>10</sup> Cyclosporine therapy is generally limited to short-term induction treatment over 30 months.<sup>13</sup>

### Surgical treatment

Surgical intervention, a proctocolectomy, can cure UC. The most common indications for surgery are acute toxic megacolon, intractable hemorrhage, poor quality of life, dysplasia, and complications from medical treatment. Surgical treatment options include proctocolectomy with ileal pouch anal anastomosis or proctocolectomy with ileostomy.

### Ileal pouch anal anastomosis

Ileal pouch anal anastomosis (IPAA) has become the operation of choice for most patients with UC.<sup>14-15</sup> This operation removes the colon (which cures UC) and preserves anorectal function. The IPAA is most often performed as a two-staged operation. An abdominal colectomy is performed, the distal rectal and proximal anal mucosa are stripped, and a pouch is constructed from the terminal ileum. While several alternatives exist for the shape of the pouch, (J, S, W, or H), the most common is the J pouch.<sup>14,16</sup> Generally, a hand-sewn anastomosis is performed, ileal pouch to anal canal. A temporary diverting loop ileostomy is created to allow fecal diversion away from the ileal pouch and ileoanal anastomosis. The loop ileostomy is closed from 2-3 months after the initial operation.

### Preoperative care

Preoperative preparation for IPAA in-




Figure 1

cludes teaching the patient about the procedure, hospital stay, long-term follow-up, and functional outcomes. This discussion should include information on living with a temporary loop ileostomy and selection of the stoma site.

Hospital stay for the uncomplicated IPAA patient is approximately 5-7 days. Patients are discharged when they tolerate a low residual diet, have adequate bowel and urinary function, have control over pain, show no signs of infection, and are able to demonstrate understanding of stomal management.

A type of stoma, the loop ileostomy, can be a challenge for patient management. Preoperative site selection is critical, because a loop ileostomy will have minimal protrusion, may create tension on the peristomal skin, and will have a high liquid effluent. Many patients who have IPAA have been steroid-dependent for months and possibly years. They may have excessive abdominal fat. Deep abdominal creases can interfere with the pouch seal. Excessive subcutaneous fat can make stomal mobilization difficult. Stoma siting is done while the patient is in a sitting position. The abdomen is evaluated for creases and folds. The belt line is considered in stoma placement. Below-the-belt stoma placement helps to conceal and support the pouch. However, if the patient has deep abdominal folds, a temporary loop stoma may not be placed below the belt line. The area chosen should have at least a 2-inch flat area that will support the pouching system. It should be located within the rectus muscle where the patient can easily see it. The proposed site is marked with a surgical marking pen and protected with a transparent dressing.

### Postoperative care

Postoperative care includes early ambulation, to support the return of bowel function and decrease the risk of complications, such as atelectasis and embolism. The Foley catheter remains in place for 4 days to prevent bladder distention in the area. A Foley catheter legband holder may be applied to prevent catheter movement and further complications. Once bowel function has returned (the presence of flatus, effluent, and bowel sounds), a clear liquid diet is started. The patient then progresses to a low residual diet. If the patient has been on long-term corticosteroids, the tapering of this medication is started.

Ileostomy output is monitored for both consistency and volume. If output is greater than 1300 cc/24 hr and liquid in consistency (normal is semi-pasty), an anti-diarrheal agent and dietary education about bulking foods are begun.

Ostomy education begins by helping the patient to learn how to empty the pouch, then progresses to changing the pouching system independently.

Follow-up in the outpatient clinic will provide the patient with continued ostomy-management education. As healing progresses, the loop stoma as well as the patient's abdomen will change in size and diameter and a convex pouching system with an extended wear barrier will usually be used.

The loop ileostomy is closed 8-12 weeks after creation of the IPAA. Prior to closure, a pouchogram is performed. It will provide information on the healing of the anastomosis. Once the pouch and anastomosis have healed, the patient is scheduled for take-down surgery. The length of hospital stay is about 5 days. Once bowel and urinary function have returned and a low residual diet is well-tolerated, the patient is discharged.

Long-term follow-up is provided after ileostomy closure to assist the patient in the use of medications and dietary modifications to reduce bowel-movement frequency and to identify symptoms suggestive of pouchitis. Stool frequency immediately after stoma closure may reach as high as 12 stools/24 hr. It will decrease as the patient starts on a regular diet.

The principal late complication after IPAA is pouchitis, which is defined as acute inflammation of the ileal pouch with mucosal ulceration. Pouchitis is characterized by episodes of a high frequency of bowel movements, passage of loose, bloodstained stools, malaise, low-grade fever, and abdominal cramping. The etiology of pouchitis is unknown. The risk of pouchitis increases over time.<sup>15</sup> Most patients with pouchitis usually respond to metronidazole in an oral dose of 750-1,500 mg/day for 7-14 days.

An examination of functional outcomes reveals the average number of stools for a patient with an IPAA is six during waking hours; one bowel movement typically occurs at night. From 20-30% of patients who have an IPAA will experience daytime incontinence. Between 41-52% of IPAA patients report spotting or soiling at night.<sup>17-18</sup>

Refinements in surgical techniques have lead surgeons to examine the role of the mucosectomy and value of a hand-sewn anastomosis versus a stapled anastomosis and the creation of an IPAA without a diverting ileostomy. There are divergent points of view on the role of the mucosectomy and type of anastomosis. The outcomes of these techniques are under review to determine the effect on the functional outcomes.

### Conclusion

The management of a patient with UC demands a team approach. This disease has no known cause or medical cure, can be socially embarrassing, and is potentially debilitating for patients and their families. It is imperative to provide education and support, as

*Continued on page 7*

**The Perioperative Management of Breast Cancer – Continued from page 1**

move the entire breast mass and a margin or normal tissue. An axillary dissection or sentinel lymph node dissection may be done at the same time.

**Nonpalpable lesions**

Two procedures are used to evaluate nonpalpable breast lesions: needle wire localization with surgical biopsy (usually lumpectomy) and stereotactic needle biopsy.

**Needle wire localization**

Needle wire localization is done in the radiology department about one to two hours before the lumpectomy procedure. A hollow-core, 20-gauge plastic needle is used. Needles are approximately 2 to 6 inches long. An appropriate length needle is selected, based on the depth of the lesion and thickness of breast tissue. The needle is inserted with mammographic, ultrasound, or stereotactic visualization guidance. A grid is superimposed on the film, allowing the radiologist to mark the coordinates of the abnormality. A local anesthetic is administered before the procedure.

After the needle is inserted, a mammogram is done to confirm accurate placement. When placement is confirmed, a flexible, fine hook wire is inserted through the hollow needle. The needle is removed and the wire stays in place. The patient goes from the radiology department to the operating room with the wire in place. The surgeon is able to feel the wire and remove it, the surrounding abnormality, and a margin of normal tissue.

It is important to inform the patient that she will be going from the radiology department to the operating room with the wire in place. All other information, support, and education about what to expect in surgical biopsy should be completed before the procedure.

A follow-up mammogram is done three months after the biopsy.<sup>3</sup>

**Stereotactic needle biopsy**

The stereotactic fine needle biopsy and stereotactic core biopsy are performed in the radiology department. Most stereotactic biopsies are done on microcalcifications seen on mammogram. Three-dimensional images are obtained by rotating the mammography unit. A computer determines the exact location of the tumor. The entire procedure takes about one hour.

The patient is positioned on a specifically designed table with an opening through which the breast is suspended. The radiologist stands under the table. Plates that are similar to those used in mammography immobilize the breast. The area is anesthetized prior to biopsy.

Several biopsies are taken from different angles. They are performed automatically by

machine with a spring-loaded device. The patient will hear a clicking sound as each biopsy is done.

The stereotactic biopsy procedure is less invasive than a surgical procedure, but women should still expect to feel mild to moderate pain afterwards due to multiple biopsy sites. Due to bruising, the nurse may suggest the patient wear a post-surgical bra that provides light support to the breast (figure 1). If the biopsies are negative for breast cancer, a follow-up mammogram is done in three months to ensure that the proper area was biopsied.<sup>4</sup>

**Patient support**

Women experience a high level of anxiety between the time that an abnormality is found and pathology results are communicated. Nurses need to provide information and much support. Information needs to be reinforced. An explanation of upcoming tests or procedures and what to expect should be given to each patient and family, supplemented by written materials.

Prior to a surgical breast biopsy or surgery, the patient needs to know:

- time and place to report.
- who will be doing the procedure.
- how long it will take.
- expected side effects.
- need for anesthesia.
- follow-up care.
- when results will be available.
- how results will be communicated.

Ambulatory nurses are responsible for coordinating these efforts and making them as seamless as possible.<sup>4</sup>

**Surgery for breast cancer**

Before treatment, a woman should discuss surgical options, such as sentinel lymph node dissection, modified radical mastectomy, axillary dissection, and modifications in radiation therapy that may minimize tissue scarring and preserve the lymphatic system.

**Sentinel lymph node dissection**

Phase three clinical trials have found that sentinel lymph node dissection is an accurate, minimally invasive way to identify breast cancer that has spread to axillary lymph nodes in women who are diagnosed with early-stage disease. Since more breast cancers are being detected early, more women will be candidates for this procedure.<sup>5</sup>

The sentinel lymph node is defined as the first node or nodes to drain the primary tu-




Figure 1. Postsurgical Bra

mor. Sentinel lymph node biopsy offers less chance of developing adverse effects, such as pain, numbness, scarring, and lymphedema, than axillary lymph node dissection. This procedure could replace axillary lymph node dissection as the standard of care.

This procedure can take from 30 minutes to 2 hours. A radioactive colloid substance, technetium, is injected around the primary tumor 30 minutes to 8 hours before surgery. If the tumor is nonpalpable, needle localization will be done to identify the tumor location. A hand-held device called a gamma probe is used to find the node(s) that have absorbed the radioactive material. The skin is marked just before surgery, then a bright blue dye is injected around the tumor before surgical preparation. This dye will travel through the lymphatic system to the sentinel node(s). The patient will receive either local or general anesthesia.

After the incision is made, the gamma probe is used to measure the sentinel node(s) before their removal. In addition, the node(s) will be stained blue.

Since sentinel lymph node biopsy is a surgical procedure, the patient may experience postoperative incisional pain. The skin may remain discolored for several days after the dye is injected. Women need to know that their urine may be green for a few hours after the procedure. Discharge instructions include care of the incision and dressing, pain control, and arm exercises.<sup>6</sup>

**Modified radical mastectomy**

Surgery for breast cancer is not new. There is evidence of breast surgery in ancient Egyptian mummies. William Halsted developed the radical mastectomy procedure in 1894. Today, the modified radical mastectomy is used. It differs from the Halsted radical mastectomy,

because the pectoral muscles and nerves are left intact. The affected breast, skin, and axillary lymph nodes are completely removed. The surgical incision extends from mid-chest to axilla. A carbon dioxide laser is now more commonly used to create the incision. The advantages of laser surgery are less blood loss and better postoperative recovery.

### Axillary dissection

Axillary dissection is done at the time of lumpectomy or mastectomy after biopsy has revealed invasive breast cancer. In some cases, a lumpectomy is done to obtain a diagnosis, then axillary dissection is performed one to two weeks later.

This procedure involves the removal of tissue within the anatomical boundaries of the axilla. The exact number of nodes will be identified by the pathologist; normally, about 6 to 15 nodes are removed. A complete dissection is not done, because it increases the risk of lymphedema.

After axillary dissection, numbness may occur in the area of surgery and along the inner arm within a rectangle from armpit to elbow. Other sensations include: numbness and tingling, burning pain, and heaviness. Most of these symptoms resolve within a year postoperatively.<sup>4</sup>

### Breast conservation surgery

Lumpectomy and axillary dissection followed by radiation therapy is as effective as mastectomy in most cases. Exceptions include the presence of breast cancer in several areas in the breast, prior radiation to the chest area, first or second trimester of pregnancy, and a history of collagen vascular disease. Other factors that would potentially prevent a patient from selecting breast conservation therapy are access to a radiation therapy center, a large tumor in a small breast that would compromise the cosmetic outcome of surgery, and a large breast size that could influence delivery of the radiation dose.<sup>7</sup>

Geography influences the choice of surgery. More breast conservation therapy is done on the East and West coasts, while more mastectomies are performed in the central part of the United States.<sup>4</sup>

Because the choice is a personal one for women, nurses need to educate them about the different treatment options that are available, procedures during treatment, the recovery process, complications, risks, and how the treatment will affect their quality of life. Nurses should also direct teaching to the patient's husband or significant other.

### Postoperative care

After mastectomy or lumpectomy with axillary dissection, the patient returns from the PACU as soon as vital signs are normal and there is no evidence of complications. The

immediate postoperative focus is on maintaining stability and comfort. Vital signs are taken frequently and dressings are assessed for bleeding. Usually one or two Jackson-Pratt drains are attached under the skin flaps to collect any fluid that would delay healing. Drainage ranges from serosanguinous to serous.

The nurse will teach the patient and family how to empty the drain and record drainage. The application of a drainage bulb holder will secure the drainage bulbs and help prevent accidental tube dislodgement. Commercially available holder with Velcro closures will simplify the procedure of emptying the bulbs. Drainage bulb holder (Dale Medical) can accommodate up to 4 bulbs and can be worn throughout the recovery period.

When the drainage is less than 25-30 cc/day, the drain will be removed. The axillary drain is usually removed in 1 to 2 weeks, but it could remain longer. The head of the bed should be elevated at least 30 degrees to help to promote drainage. The affected arm is elevated on a pillow while the patient is awake. Keeping the arm elevated will help to promote drainage until collateral lymph channels develop. The patient should have pneumatic compression on her lower extremities while bedridden. Postoperative activities, such as coughing, turning, deep breathing, and incentive spirometry, are followed until the patient is mobile. An abdominal binder with Velcro closure may be prescribed to help encourage coughing, turning, and deep breathing while splinting the diaphragm for the patient. Basic comfort measures, such as turning and repositioning, are done as needed. Pain needs to be assessed and managed on a regular basis.<sup>8</sup>

Nurses can refer women to the American Cancer Society's Reach to Recovery program before surgery or postoperatively. A Reach to Recovery volunteer who has had the same surgery will serve as her peer counselor.

Women are instructed that a permanent prosthesis can be purchased several weeks after surgery, when the mastectomy incision has healed. The prosthesis is a breast form that fits in or is attached to the chest wall, a bra, bathing suit, or lingerie. The store where a patient purchases the prosthesis should employ a trained fitter. Medicare and most insurance will reimburse the cost of a prosthesis with a note from the doctor.

### Care of incisions after discharge

Usually the surgeon will remove the dressings before the patient is discharged. The patient should be instructed how to care for the incision. If she is allowed to shower, she will use a Jackson-Pratt drain pouch to keep the drain from accidentally pulling out. During bathing, the drain bulb will float in the tub. Advise the patient not to panic if the drain pulls out. Instruct her to place a gauze dressing over

the site and call her surgeon's office for more instructions.

Advise the patient to call the surgeon immediately if:

- her temperature is over 101°F.
- the incision starts to bleed. If bleeding is persistent, she must go directly to the emergency room.
- the incision suddenly becomes hot, red, tender, or drains pus.<sup>9</sup>

Postoperative arm exercises will help to develop collateral lymphatic channels. If the patient remains overnight in hospital, she will have a visit from physical therapy to review the exercises. If she goes home on the day of the procedure, the nurse will review the exercises with her and her family.

Exercises that use the hand and forearm are introduced on the day of surgery and first postoperative day. Day one exercises continue on day two, but range of motion is slightly increased. The patient is instructed to touch the opposite shoulder, touch the opposite knee, comb her hair, and reach for the small of the back with the affected arm. On day three, range of motion exercises are further increased. The goal is for the patient to be able to fully extend her arm after several days. Exercises from days one and two continue, and other exercises, such as pendulum, jump rope, back scratch, wall climbing, clasp, reach, and spread, and overhead pulley exercises are added. If the patient has difficulty in progressing, she is referred to outpatient physical therapy.

### Hand and arm care

To avoid cuts, scratches, irritations, and burns, advise patients to:

- use cuticle cream instead of scissors when cutting nails.
- wear a thimble for sewing.
- use non-irritating deodorants or perfumes.
- use insect repellent when outdoors.
- use rubber gloves for washing dishes and handling harsh cleansers.
- use an electric razor for underarm shaving.
- wear gloves and long sleeves for gardening.
- use a long, padded glove when reaching into a hot oven or barbecue.
- use sunscreen to prevent sunburn.

Web sites

**American Cancer Society: [www.cancer.org](http://www.cancer.org)**

This site describes the different types of cancer, while discussing methods of prevention and treatment. It includes recent news about clinical trials and research. It also contains statistical information.

**Breast Cancer Info.com: [www.breastcancerinfo.com](http://www.breastcancerinfo.com)**

This site contains information about upcoming events related to breast cancer, and it has updates on the latest professional conferences. It has information about treatments and research studies.

**Cancer Information Network: [www.cancernetwork.com](http://www.cancernetwork.com)**

This site provides information for health-care professionals, patients, and the families of patients. It includes general descriptions of the different types of patient information.

**Cancer Survivors Network: [www.cancersurvivorsnetwork.org](http://www.cancersurvivorsnetwork.org)**

Run by the American Cancer Society, this network is a means by which cancer survivors can communicate. The site contains personal stories, and it addresses the future for cancer survivors.

**Susan G. Komen Foundation: [www.komen.org](http://www.komen.org)**

This site has detailed information on breast health services, including the education, research, clinical, and breast cancer screening programs of the Susan G. Komen Breast Cancer Foundation.

**Living Beyond Breast Cancer: [www.lbbc.org](http://www.lbbc.org)**

This site includes educational materials and updates about special events relating to breast cancer. It also has links related specifically to breast cancer.

**National Cancer Institute (NCI): [www.nci.nih.gov](http://www.nci.nih.gov)**

This site contains announcements about upcoming events and research programs. It provides information about managing both the physical and emotional effects of cancer.

**Y-ME Breast Cancer Organization: [www.y-me.org](http://www.y-me.org)**

This site contains general breast cancer information. There is also a special section addressing breast cancer in men.

To minimize binding or constricting, advise patients to:

- avoid tight jewelry or clothing
- avoid blood drawing, injections, or IVs on the side of surgery
- carry a purse on the unaffected side
- ask that blood pressure be taken on the unaffected arm<sup>9</sup>

**Postoperative complications**

About one-third of women who undergo axillary lymph node dissection develop lymphedema in the affected arm. Surgery, radiation therapy, and infection are all contributing factors in its development.

Lymphedema occurs when lymph nodes and vessels are damaged during treatment. Lymphatic flow is impeded, then fluid accumulates in the soft tissue around the axillary dissection site and may extend into the arm. Once lymphedema develops, it can be managed but cannot be cured. Lymphedema may develop weeks, months, or years after the initial surgery.

Nurses should obtain a baseline measurement of the affected arm before surgery, six weeks postoperatively, and any time that swelling occurs. Patients should be instructed to report a difference between the two arms, such as a heaviness or a change in how clothes fit.

Interventions for all women who experience lymphedema include elevation, reduction in salt intake, massage, mild exercise, avoidance of heat or trauma, and elastic support. An intermittent sequential pneumatic compression sleeve is needed for moderate to severe lymphedema.

Physical discomfort is only one component of the lymphedema experience. Women may need ongoing support to cope with this chronic problem. A multidisciplinary team should be available, including physical therapy, occupational therapy, social work, and counselors.<sup>10</sup>

**Breast reconstruction**

Reconstruction can be done at the time of mastectomy or any time afterwards. Options for breast reconstruction should be discussed before surgery. There are two types of breast reconstruction: breast implant or tram flap. Both procedures are done by a plastic surgeon.

**Breast implants**

The saline-filled breast implant is used to create a breast mound that looks similar to a natural breast. The saline-filled shell is implanted in the chest muscle. A nipple can be reconstructed after the implant procedure is completed. The silicone-filled breast implant is only available through controlled clinical trials because of safety concerns.

A tissue expander is used if there is not enough skin and chest muscle to cover an implant. A tissue expander is an empty saline shell that is injected with saline once weekly through a metal port. This procedure gradually stretches the muscle and tissue. When expansion is complete, a permanent breast implant is placed.

Women need to be told that a saline implant is similar to a natural breast but does not have exactly the same look or feel. Surgery is sometimes needed to match the remaining breast with the breast implant.

**Tram-flaps**

The tram flap (transverse rectus abdominis myocutaneous flap) involves abdominal surgery. A breast is reconstructed

from abdominal muscle, fat, and skin. It feels more like a normal breast, because the patient's own tissue is used. A flap of skin and muscle from the lower abdomen is attached to one of the rectus muscles in the abdomen. A tunnel is made under the skin from the abdomen to the mastectomy site. The skin flap, fat, and muscle are pulled through the mastectomy incision, then formed into the shape of a breast.

The patient has a nasogastric tube postoperatively. Several Jackson-Pratt drains are in place. Pain control is often managed with PCA morphine for the first few postoperative days. The patient is typically hospitalized from 3 to 5 days. Arm exercises are usually delayed until the flap has healed. Coughing, turning, deep breathing, incentive spirometry, and pneumatic compression stockings are used postoperatively, until the patient is ambulatory.

A nipple and areola will be created after the initial surgery has healed. The formed breast may need further adjustments. A post-surgical bra that provides light compression may be applied to hold surgical dressings in place without the need for tape or ace bandages. Overall, the cosmetic outcome of tram-flap reconstruction is very good.

Breast cancer is a challenging disease for patients and their families. Nurses must stay informed about new surgical techniques for diagnosing or treating breast cancer in order to continue to provide information and support throughout the perioperative experience.

*References*

1. A Cancer Journal for Clinicians. Cancer Statistics 2001. Jan/Feb 2001, 51(1):16 and 23.
2. Hunter MA, MD. *The Little Book of Breast Cancer: A Self-Teaching Guide*. Cancer Info Press: Mercer Island, WA. 1998. pp. 17.
3. Kelly P, Winslow E. Needle wire localization for nonpalpable breast lesions: sensations, anxiety levels and informational needs. *Oncology Nursing Forum*. 1996; 23 (4): 639-645.
4. Tish M. *Cancer Nursing Breast Cancers*. McCorkle R, Grant M, Frank-Stromborg M, Baird S. (eds.). WB Saunders: Philadelphia. p. 560-569.
5. Zack E. Sentinel lymph node biopsy in breast cancer: Scientific Rationale and Patient Care. *Oncology Nursing Forum*. 2001; 28 (6): 997-1005.
6. Baron R. Sentinel lymph node biopsy in breast cancer and the role of the oncology nurse. *Clinical Journal of Oncology Nursing*. 1999; 3(1): 17-22.
7. Osteen R. Breast Cancer. In: Lenhard JR, Osteen RT, Gansler MP (eds.). *American Cancer Society Clinical Oncology*. Atlanta, GA. 2001. pp. 260-262.
8. Ignatavicius DD, Worknam LM, Mishler MA. Interventions for the clients with breast disorders. *Medical Surgical Nursing: A Nursing Process Approach*. WB Saunders: Philadelphia, 1995.
9. Rummel M, Deming P. Patient guide to breast surgery: Pennsylvania Hospital: Joan Karnell Cancer Center. 2001. Philadelphia.
10. Lockhart JS. Case in point lymphedema. *Oncology Nursing Forum*. 1999; 26 (3).



**Cathy Fortenbaugh,** AOCN, CSN, MSN, BSC, is an Oncology Clinical Nurse Specialist for a 42-bed medical/surgical oncology unit at Pennsylvania Hospital, Philadelphia, PA. She is Chair of the Nursing Research Council and a member of the Hospital Research Review Committee, Hospital Ethics Committee, and University of Pennsylvania Health System Nursing Research Committee. Cathy acts as an oncology consultant to hospital staff, peers and the community.

**Management of Patients with Inflammatory Bowel Disease – Continued from page 3**

treatment focuses on either medical or surgical intervention.

Team members must include a gastroenterologist, IBD surgeon, dietitian, social worker, and certified wound, ostomy, and continence nurse. Collaboration by the team members should assure good outcomes for the patient living with UC.

**References**

- Hanauer SB. Inflammatory bowel disease. In: Bennett JC, Plum F (eds.). Cecil Textbook of Medicine. 20<sup>th</sup> ed. Saunders: Philadelphia, 1996, pp. 707-715.
- Sandler, RS, Eisen GM. Epidemiology of inflammatory bowel disease. In: Kirsner JB (ed.). Inflammatory Bowel Disease. 5<sup>th</sup> ed. Saunders: Philadelphia, 2000, pp. 89-112.
- Brant SR, Cho JH. Inflammatory bowel disease genetics. In: Bayless TM, Hanauer SB (eds.). Advanced Therapy of Inflammatory Bowel Disease. Becker: Hamilton, 2001, pp. 523-526.
- Mayer L. Current concepts of inflammatory bowel disease etiology and pathogenesis. In: Kirsner JB (ed.). Inflammatory Bowel Disease. 5<sup>th</sup> ed. Saunders: Philadelphia, 2000, pp. 280-296.
- Ahnen DJ. Gastrointestinal malignancies in inflammatory bowel disease. In: Kirsner JB (ed.). Inflammatory Bowel Disease. 5<sup>th</sup> ed. Saunders: Philadelphia, 2000, pp. 379-396.
- Miner PB. Clinical features, course, laboratory findings, and complications in ulcerative colitis. In: Kirsner JB (ed.). Inflammatory Bowel Disease. 5<sup>th</sup> ed. Saunders: Philadelphia, 2000, pp. 299-314.
- Saxon A, Shanahan F, Landers C, et al: A distinct subset of antineutrophil cytoplasmic antibodies is associated with inflammatory bowel disease. *J Allergy Clin Immunol* 1990;86:202-210.
- Svartz N: Sulfasalazine: II. Some notes on the discovery and development of salazopyrin. *Am J Gastroenterol* 1988;83:497-503.
- Greenfield SM, Punched NA, Teare JP, et al. The mode of action of the aminosalicylates in inflammatory bowel disease. *Aliment Pharmacol Ther* 1993;7:369-383.
- Hanauer SB. Medical therapy for ulcerative colitis. In: Kirsner JB (ed.). Inflammatory Bowel Disease. 5<sup>th</sup> ed. Saunders: Philadelphia, 2000, pp. 529-556.
- Kornbluth AA, Salomon P, Sacks Hs, et al. Meta-analysis of the effectiveness of current drug therapy of ulcerative therapy. *J Clin Gastroenterol* 1993;9:215-218.

- Olmstead J. Evaluation and management of the patient with ulcerative colitis. *Gastroenterology Nursing* 1998;21:176-182.
- Kornbluth A, Present DH, Lichtiger S, et al.: Cyclosporin for severe ulcerative colitis: a user's guide. *Am J Gastroenterol* 1997;92:1424-1428.
- Remzi FH, Fazio VW. Ileoanal pouch anastomosis. In: Bayless TM, Hanauer SB (eds.). Advanced Therapy of Inflammatory Bowel Disease. Becker: Hamilton, 2001, pp. 198-202.
- Dozios RR, Kelly KA. The surgical management of ulcerative colitis. In: Kirsner JB (ed.). Inflammatory Bowel Disease. 5<sup>th</sup> ed. Saunders: Philadelphia, 2000, pp. 626-657.
- Michelassi F, Hurst R. Restorative proctocolectomy with j-pouch ileoanal anastomosis. *Arch Surg* 2000;135:347-535.
- Gray M, Colwell JC. Evidence-based report card: what functional outcomes and complications should be taught to the patient with ulcerative colitis or familial adenomatous polyposis who undergoes ileal pouch anal anastomosis. *J Wound, Ostomy and Continence Nursing* 2001;28, in press.
- Meagher AP, Farouk R, Dozios RR, Kelly KA, Pemberton JH. J ileal pouch anal anastomosis for chronic ulcerative colitis: complications and long term outcomes in 1310 patients.



**Janice C. Colwell,** RN,MS,CWOCN, is a specialist, educator, and consultant in wound, ostomy, and continence nursing. Based in Chicago, Janice plays an active role in the Wound, Ostomy, and Continence Nurses Society, where she is Chair of the Communications Committee and Ostomy Section Editor of the WOCN journal.

*Perspectives*, a quarterly newsletter focusing on post-operative recovery strategies, is distributed free-of-charge to health professionals. *Perspectives* is published by Saxe Healthcare Communications and is funded through an education grant from Dale Medical Products Inc. The newsletter's objective is to provide nurses and other health professionals with timely and relevant information on postoperative recovery strategies, focusing on the continuum of care from operating room to recovery room, ward, or home.

The opinions expressed in *Perspectives* are those of the authors and not necessarily of the editorial staff, Cross Country University, or Dale Medical Products Inc. The publisher, Cross Country University and Dale Medical Corp. disclaim any responsibility or liability for such material.

We welcome opinions and subscription requests from our readers. When appropriate, letters to the editors will be published in future issues.

Please direct your correspondence to:  
**Saxe Healthcare Communications**  
**P.O. Box 1282, Burlington, VT 05402**  
**Fax: (802) 872-7558**



Cross Country University is an accredited provider of continuing education in nursing by the American Nurses Credentialing Commission on accreditation.

After reading this educational offering, the reader should be able to:

- Review the role of surgery in the treatment of breast cancer.
- Differentiate between the five types of biopsies used to diagnose breast cancer.
- Discuss breast-conservation therapy and the advantages and disadvantages of each therapy.
- Describe the perioperative nursing interventions for the woman who has surgery for breast cancer.
- Discuss two different types of reconstruction options for women who have a mastectomy.

**To receive continuing education credit, simply do the following:**

- Read the educational offering.
- Complete the post-test for the educational offering. Mark an X next to the correct answer. (You may make copies of the answer form.)
- Complete the learner evaluation.
- Mail, fax, or send on-line the completed learner evaluation and post-test to the address below.
- 1.0 contact hours for nurses are awarded by Cross Country University, the Education and Training Division of Cross Country Inc., which is accredited as a provider of continuing education in nursing by the American Nurses Credentialing Center's Commission on Accreditation. Cross Country University is an approved provider with the Iowa Board Of Nursing, approved provider #328. This course is offered for 1.0 contact hours. Cross Country University is approved by the California Board of Registered Nursing, Provider #CEP 13345, for 1.0 contact hours.
- To earn 1.0 contact hours of continuing education, you must achieve a score of 75% or more. If you do not pass the test, you may take it again one time.
- Your results will be sent within four weeks after the form is received.
- The administrative fee has been waived through an educational grant from Dale Medical Products, Inc.
- Answer forms must be postmarked by April 30, 2006, 12:00 midnight.

Name \_\_\_\_\_  
 Credentials \_\_\_\_\_  
 Position/title \_\_\_\_\_  
 Address \_\_\_\_\_  
 City \_\_\_\_\_ State \_\_\_\_ Zip \_\_\_\_\_  
 Phone \_\_\_\_\_  
 Fax \_\_\_\_\_  
 License #: \_\_\_\_\_  
 \* Soc. Sec. No. \_\_\_\_\_  
 E-mail \_\_\_\_\_  
 \* required for processing

Mail to: Cross Country University  
 6551 Park of Commerce Blvd. N.W., Suite 200  
 Boca Raton, FL 33487-8218

or: Fax: (561) 988-6301  
 www.perspectivesinnursing.org

- A needle localization procedure is done:**
  - for palpable lesions seen on mammogram
  - for nonpalpable lesions seen on mammogram
  - for all women who have breast-cancer surgery
  - for microcalcifications that a woman can feel
- The sentinel node is defined as:**
  - the first node or nodes to which cancer has spread
  - the most distant nodes in the chain
  - the most proximal nodes in the chain
  - the first node or nodes that drain the primary tumor
- Lymphedema occurs in:**
  - all women postoperatively
  - almost no women due to advances in surgical techniques
  - one-third of women who have axillary lymph node dissection
  - three-fourths of women who have axillary lymph node dissection

- An incisional breast biopsy is indicated:**
  - to obtain a tissue diagnosis in a large tumor
  - to remove the entire tumor and a margin of normal tissue
  - to conserve as much breast tissue as possible
  - after breast cancer has reoccurred at the chest wall
- Immediate postoperative arm exercises are NOT indicated for which of the following procedures:**
  - mastectomy
  - lumpectomy with axillary lymph node dissection
  - tram-flap reconstruction
  - sentinel lymph node biopsy
- At what point will the Jackson-Pratt drain be removed after axillary lymph node dissection?**
  - prior to discharge
  - postoperative day one
  - when the drainage is less than 60-100 cc a day
  - when the drainage is less than 25-30 cc a day

- The American Cancer Society's Reach to Recovery Program is:**
  - a home physical therapy program to reinforce arm exercise
  - a chronic lymphedema management program
  - a peer counselor program focusing on education and support
  - a hospital-based support group
- The diagnosis of breast cancer is definitively made by:**
  - mammogram
  - breast self-examination
  - evaluation of breast tissue by a pathologist
  - radiologic evaluation
- After mastectomy, pain is:**
  - managed exclusively by epidural
  - monitored closely
  - managed exclusively by PCA morphine
  - assessed and managed on a regular basis

**Mark your answers with an X in the box next to the correct answer**

1	A B C D E	3	A B C D E	5	A B C D E	7	A B C D E	9	A B C D E
	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
2	A B C D E	4	A B C D E	6	A B C D E	8	A B C D E		
	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>		

**Participant's Evaluation**

1. What is the highest degree you have earned?      1. Diploma      2. Associate      3. Bachelor's      4. Master's      5. Doctorate  
 Using 1 =Strongly disagree to 6= Strongly agree rating scale, please circle the number that best reflects the extent of your agreement to each statement.

**Strongly Disagree**

**Strongly Agree**

2. Indicate to what degree you met the objectives for this program:

■ Review the role of surgery in the treatment of breast cancer.	1	2	3	4	5	6
■ Differentiate between the five types of biopsies used to diagnose breast cancer.	1	2	3	4	5	6
■ Discuss breast-conservation therapy and the advantages and disadvantages of each therapy.	1	2	3	4	5	6
■ Describe the perioperative nursing interventions for the woman who has surgery for breast cancer.	1	2	3	4	5	6
■ Discuss two different types of reconstruction options for women who have a mastectomy.	1	2	3	4	5	6

- Have you used home study in the past?     Yes     No    \_\_\_\_\_
- How many home-study courses do you typically use per year?    \_\_\_\_\_
- What is your preferred format?     video     audio-cassette  
     written     combination    \_\_\_\_\_
- What other areas would you like to cover through home study?    \_\_\_\_\_

For Iowa nurses, you may submit the evaluation to Iowa Board of Nursing.

Mail to: Cross Country University, 6551 Park of Commerce Blvd. N.W., Suite 200, Boca Raton, FL 33487-8218• or Fax: (561) 988-6301  
 E-mail: [perspectivesinnursing.org](mailto:perspectivesinnursing.org)